

Report on field quality in the main LHC dipoles January-February 2005

E. Todesco, AT-MAS-MA

This report gives data relative to field quality measured in collared coils and cold masses during the period January 1– February 28 2005, warm-to-cold correlations, comparison to beam dynamics targets, and status of the holding points. Updated graphs can be found in the field quality observatory <http://lhc-div-mms.web.cern.ch/lhc-div-mms/MMSPAGES/MA/Obs.html>.

EDMS n. 577313

The dashboard

- Available measurements:
 - 730 collared coils and 676 cold masses at room temperature,
 - 133 dipoles plus 8 half dipoles (one aperture only) for a total of 137 equivalent dipoles at 1.9 K¹.
- In these two months we had:
 - 70 new collared coils (20 from Firm1, 19 from Firm2 and 31 from Firm3) measured at room temperature,
 - 4 dipoles measured at 1.9 K¹.

What's new (not much)

- **Production rate** is at 35 collared coils per month. The rate is 2.3, 2.2 and 3.6 collared coils per week in Firm1, Firm2 and Firm3, respectively.
- **Length of feedback loop:** The minimal delay between collared coil magnetic measurements and magnetic measurements at 1.9 K is 50 days (obtained for magnet 1136). The last magnet measured at 1.9 K is the 605th, and since then another 125 collared coils have been produced.
- **Change of target for systematic b3:** A revision of the target range for the systematic b3 at injection has been agreed between AT-MAS and AB-ABP. The window has been reduced from [-10.5,10.5] units to [-7,7] units. This corresponds to cut the lower half of the target range on the room temperature measurements (see also Figs. 13 and 14, pg. 9). This reduction has been asked to have a more realistic bound on the beta beating induced by sextupolar spool pieces misalignment at injection. This has allowed to increasing tolerances for spread of integrated field gradient in the quadrupoles, which is critical.
- **No trends:** Transfer function and multipoles are stable along the production of the last two months. Correlations are stable.
- **No assembly faults:** No assembly faults detected in these two months.

¹ These numbers refers to measurements of either magnets or single apertures available in AT-MTM Oracle database at the time of distribution of the report.

CONTENTS

PART I: MEASURED MAGNETS AND ASSEMBLY DATA.....	pg. 3
PART II: MEASUREMENTS VERSUS BEAM DYNAMICS TARGETS.....	pg. 4
2.1 Summary of systematics components.....	pg. 4
2.2 Summary of random components.....	pg. 5
PART III: TRENDS IN FIELD QUALITY.....	pg. 6
3.1 Trends in bending strength.....	pg. 6
3.2 Trends in normal odd multipoles.....	pg. 9
3.3 Trends in normal even multipoles.....	pg. 11
3.4 Trends in skew multipoles.....	pg. 14
3.5 Trends in systematic differences between Firms.....	pg. 15
3.6 Trends in correlations to measurements at 1.9 K.....	pg. 16
PART IV: QUALITY CONTROL.....	pg. 18
4.1 Holding point results.....	pg. 18
4.2 Coil waviness.....	pg. 19

The new format of the report

We remind the reader the most important features of the report.

- The first section deals with the number of measured magnets in the last two months and the assembly data (X-section type and shim size).
- In the second section we have the summary of the measured field quality of all collared coils versus beam dynamics targets. This gives a quick overview of the best guess for the status of field quality versus beam dynamics.
- The third section is devoted to trends in field quality.
 - The trend plots show multipole moving averages for each manufacturer versus the magnet progressive number². Each marker is the average of 5 measurements:
 - the collared coil characterized by the progressive number in the horizontal axis
 - the two collared coils previously produced by the same Firm
 - the two collared coils produced afterwards by the same Firm
 - We always give plots for the collared coil measurements, except the case of bending strength where also cold masses measurements are adding important information. When comparing these cold masses to collared coils, one has to take into account that usually the last 60 collared coils have not yet become cold masses, and therefore a different pattern has to be expected in the end of the plot (see Figs. 9-10, and 11-12).
 - We give the reduction to nominal shims only for b3. Now shims are nearly always nominal.
 - Correlations are not presented in the standard plot 'warm-vs-cold', but rather as a trend plot of the offset between warm and cold vs the magnet progressive number. In this way we can visualize trends in correlations and the type of sampling that is being carried out at 1.9 K.
 - From this report on, all plots give integral values (i.e. including contribution of coil heads). This gives in some cases an offset in the y scale with respect to plots of the previous reports, corresponding to the head contribution.
- The final Section is devoted to field quality used to detect a faulty assembly procedure.

² We recall the definition of magnet progressive number, used as abscissa axis in most of our trend plots: it is a number running from 1 to 1232 which is associated to each magnet, according to the date of the first magnetic measurement at room temperature.

PART I: MEASURED MAGNETS AND ASSEMBLY DATA

- 70 new collared coils have been measured (collared coils 661st to 730th).
 - 20 of Firm1 (1127,1141,1154,1165,1202-6,1208-16,1218-9)
 - 19 of Firm2 (2152-2170)
 - 31 of Firm3 (3242,3312,3315-22,3324-45)




Fig. 1: Number of magnets measured at CERN at 1.9 K and at the manufacturers at room temperature at different stages of assembly procedure

- Cross-section: collared coils have X-section 3.
- Shims are nominal in all Firms, except one case in Firm3 (see Fig. 2).




Fig. 2: Thickness of the polar shims used in the collared coils

PART II: MEASUREMENTS VERSUS BEAM DYNAMICS TARGETS

2.1 Summary of systematic components

- Best estimates of skew and even normal systematic components are given in Fig. 3. All the multipoles are within specifications. Details on trends are given in Part III.




Fig. 3: Best estimate for systematic skew multipoles and even normal multipoles (markers) versus beam dynamics limits (red line).

- Best estimates for systematic odd normal multipoles are shown in Fig. 4. The revision of the targets on systematic b_3 has reduced the minimal acceptable systematic b_3 in the collared coil from -8.2 units (see previous report) to -4.5 units. In the left part, raw data are plotted. This gives the actual situation for global values relative to all manufactured collared coils, which are slowly moving towards optimal ranges: b_3 is within target and b_5 is larger than the upper target of 0.18 units.
- In the right part of Fig. 4, data are separated according to the three cross-sections (34 collared coils have cross-section 1, 147 have cross-section 2, 548 have cross-section 3, plus one hybrid 1-2). With cross-section 3, b_3 in the collared coil is 1.3 units below the upper limit (i.e., 1.9 units at high field), and also b_5 is within targets, just below the edge of the upper limit (i.e., 1.08 units at injection). Finally, b_7 in the collared coil is 0.27 units larger than the limits (i.e. 0.34 units at injection).




Fig. 4: Best estimate for systematic odd normal multipoles (markers) versus beam dynamics limits (red line). Raw data (left) and data separated according to different cross-sections (right).

2.2 Summary of random components

- We evaluate the standard deviation of the bending strength and multipoles for each Firm and for all magnets (see Fig. 5). We analyse only magnets with cross-section 3 (548 collared coils). Standard deviation of multipoles in collared coil are divided by 1.18 to give the best estimate of the random due to geometric in the cold mass, and compared to the target for the beam dynamics (**whose budget includes the random components induced at 1.9 K**). All values are well within targets, with the exception of the main field in straight part B; please note that the relevant constraint for beam dynamics is only on the bending strength BdL, which is within targets.




Fig. 5: Random component in the measured collared coils and rescaled to cold mass values, cross-section 3 only compared to targets for random at 1.9 K.

- The analysis of the spread per sector is being reviewed on the basis of the installation constraints. A complete picture will be available in the next report

PART III: TRENDS IN FIELD QUALITY

3.1 Trends in bending strength

3.1.1 Trends in magnetic length

- Magnetic length of the collared coils is extremely stable in all Firms since magnet progressive number 100 (see Fig. 7). Magnetic length in Firm1 is 5 units higher than in Firm2 and Firm3.




Fig. 7: Magnetic length of the measured collared coils, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm).

- Magnetic length of **cold masses** is also extremely stable in all Firms since magnet progressive number 100 (see Fig. 8). When iron laminations are added, magnetic length in Firm3 is getting smaller than in Firm1 and 2. The net result is that there are around 8 units of difference between Firm1 and Firm3, with Firm2 in between.




Fig. 8: Magnetic length of the measured **cold masses**, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm).

3.1.2 Trends in transfer function

- Transfer function in collared coils 660th to 730th is rather stable in all Firms. All Firms are within ± 15 units (see Fig. 9).




Fig. 9: Transfer function of the measured collared coils, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm).

- The systematic difference in the transfer function between Firms observed in collared coils is confirmed, but reduced of around 20% (i.e., the iron yoke contribution), in **cold mass** data (see Fig. 10).




Fig. 10: Transfer function of the measured **cold masses**, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm).

3.1.3 Trends in integrated transfer function

- The integrated transfer function shows a spread between Firms of at most ± 5 units in recent production (see Fig. 11).




Fig. 11: Integrated transfer function of the measured collared coils, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm).

- In the **cold masses** data the spread of the integrated transfer function between Firms is reduced by 20% (see Fig. 12).




Fig. 12: Integrated transfer function of the measured cold masses, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm).

3.2 Trends in odd normal multipoles

- Average b_3 in all Firms in the collared coils is between -1 and -3 units. The recent production shows a small spread between firms. The negative trend of Firm2 around magnet progressive number $360^{\text{th}}-540^{\text{th}}$ is over. We remind the reader that the peak in Firm1 collared coils around magnet progressive number 430 is due to the additional cross-section 2 magnets.
- Please note the in the Figures we implemented the new target on systematic b_3 at injection $[-7,7]$ instead of the previous value of $[-10.5,10.5]$.




Fig. 13: Average b_3 in straight part of the collared coils, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.




Fig. 14: Average b_3 in straight part of the collared coils, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles. Data reduced to nominal shims³.

³ This plot is very similar to the plot of Fig. 13 since shims are nominal for most of the production

- The positive trend of b_5 in Firm2 observed between 380th and 520th has disappeared. Measured b_5 in Firm2 is decreasing, reaching the lower limit (see Fig. 15).
- Average b_5 in Firm3 is also slightly decreasing, but well within targets.
- Average b_5 in Firm1 is around 0.3 units, i.e. around 1 unit higher than in Firm2 and Firm3.
- Systematic b_5 in the recent production of these two months has decreased and is around -0.5 units, i.e. within target.




Fig. 15: Average b_5 in straight part of the collared coils, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.

- Normal 14th pole b_7 is stable.




Fig. 16: Average b_7 in straight part of the collared coils, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.

3.3 Trends in even normal multipoles

For each multipole being subject to beam dynamics specifications, we present two separated plots for the systematic per aperture, plus a plot of the systematic per beam, i.e. the average of both apertures (that should be zero due to the two-in-one symmetry).

3.3.1 Trends in normal quadrupole

- The systematic per aperture is in the upper (lower for aperture 2) part of the target range (see Figs. 17 and 18).




Fig. 17: Average b2 in straight part of the collared coils (aperture 1), separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.




Fig. 18: Average b2 in straight part of the collared coils (aperture 2), separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.

- The systematic normal quadrupole per beam is within specifications (see Fig. 19).




Fig. 19: Average b_2 in straight part of the collared coils (average of the apertures), separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.

3.3.2 Trends in normal octupole

- The systematic per aperture is within specifications in both apertures (see Figs. 20 and 21).
- The systematic per beam is also within specifications (see Fig. 22).




Fig. 20: Average b_4 in straight part of the collared coils (aperture 1), separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.




Fig. 21: Average b_4 in straight part of the collared coils (aperture 2), separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.




Fig. 22: Average b_4 in straight part of the collared coils (average of the apertures), separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.

3.4 Trends in skew multipoles

- Skew quadrupole a_2 is well within targets, and no trends are observed (see Fig. 23). Firm3 has a systematic component of 0.5 to 1 unit since magnet 200th, whereas Firm2 and Firm1 are well centred around zero.




Fig. 23: Average a_2 in straight part of the collared coils, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.

- Skew sextupole a_3 is well within targets (see Fig. 24). There is a positive systematic component in Firm3 (around 0.5 units), and a slightly negative component (around 0.25 units) in Firm1 and Firm2. Indeed, beam dynamics targets are very loose, and therefore there is no concern on this multipole.




Fig. 24: Average a_3 in straight part of the collared coils, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.

- Skew octupole a_4 is within the tight beam dynamics targets in Firm1 (see Fig. 25).
- The strong systematic component in Firm2 is oscillating between 0 and 0.7 units. Recent production of Firm2 features an a_4 closer to zero.
- The strong systematic a_4 in Firm2 is partially compensated by negative values in the production of Firm3.




Fig. 25: Average a_4 in straight part of the collared coils, separated per Firm (each dot is average of 5 consecutive magnets of the same Firm), and beam dynamics targets for the systematic (red lines) based on correlations with 134 cryodipoles.

3.5 Trends in systematic differences between Firms

The more relevant signature of Firms is in b_7 and a_3 .

- Normal 14th pole: b_7 at Firm2 is 0.5 units lower than Firm3 and Firm1 (see Fig. 16). This difference is three times the natural sigma within the same manufacturer measured in cross-section 3. Firm2 is within targets, whereas both Firm1 and Firm3 are outside.
- Skew sextupole a_3 : Firm3 has a systematic a_3 of 0.5 units, against -0.5 units in Firm1-2 (see Fig. 24). This difference is three times the natural sigma within the same manufacturer. All Firms are within targets.

We observe a small systematic difference between Firms (from one to two times the natural sigma within the same manufacturer) in the following cases:

- Normal decapole b_5 : Firm1 has a systematic b_5 of 1 unit larger than Firm2-3. This difference is two times the natural sigma within the same manufacturer (see Fig. 15). Firm2-3 are within targets, whereas Firm1 is outside.
- Skew octupole a_4 : Firm2 has a systematic a_4 of 0.4 units, against -0.03 and -0.05 units in Firm3 and Firm1, respectively (see Fig. 25). This difference is equal to the natural sigma within the same manufacturer. Firm1 and Firm3 are within targets, whereas Firm2 is outside.

Systematic differences between Firms are small or negligible in a_2 , b_2 , b_3 and b_4 .

3.6 Trends in correlations to measurements at 1.9 K

We give plots of the offsets between the values measured at injection field (or high field) at 1.9 K, without beam screen, and the cold mass at room temperature⁴. The offsets are given versus the magnet progressive number. This gives a hint on the type of sampling of the production that is being carried out with the measurement at 1.9 K. The last magnet measured at 1.9 K is collared coil 605th, thus implying a delay of 125 collared coils with respect to the last manufactured collared coil (i.e. the 730th), which corresponds to three months of production, i.e. around 3/4 of octant.

- Trend plots for the offsets relative to the integrated transfer function are given in Figs. 26 and 27, at injection and at high field, respectively. In both cases no trends are visible after collared coil 100th.




Fig. 26: Difference for the integrated transfer function between measured values at 1.9 K, injection field, and **cold mass at r.t.** along the magnet production.




Fig. 27: Difference for the integrated transfer function between measured values at 1.9 K, high field, and **cold mass at r.t.** along the magnet production.

⁴ Please note that in previous reports we gave the difference with respect to the collared coil.

- We present data relative to b3-injection and b3-high field in Figs. 28 and 29. Please note the enlarged scale with respect to b3 plots in Figs. 13 and 14. Offsets are stable, and fall within a range of ± 0.7 units at injection. At high field the offset is very stable (within ± 0.5 units).




Fig. 28: Difference for the b3 between measured values at 1.9 K, injection field, and **cold mass at r.t.**, along the magnet production.




Fig. 29: Difference for the b3 between measured values at 1.9 K, high field, and **cold mass at r.t.**, along the magnet production.

- Trends for the b5 and b7 offsets between injection and collared coil straight part are given in Fig. 30 and 31. The situation is stable in both cases.




Fig. 30: Difference for the b5 between measured values at 1.9 K, injection field, and **cold mass at r.t.**, along the magnet production.




Fig. 31: Difference for the b7 between measured values at 1.9 K, injection field, and **cold mass at r.t.**, along the magnet production.

PART IV: QUALITY CONTROL

4.1 Holding point results

We had no few cases of relevant field anomalies

A summary of the magnets de-collared for anomalies in the magnetic field over all the production is given in Table I. The total number of found defects is 13 over 730 collared coils, i.e. 1.8%. A large fraction of these defects (8 over 13) has been found in collared coil 300th to 400th (see Fig. 32). The situation is improving in the more recent production.




Fig. 32: Total number of defects found with magnetic measurements versus magnet progressive number.

Table I: Summary of magnets decollared on the basis of anomalies in magnetic field.

Bad assembly cases				
Magnet	Measured on	Analysis	Opened on	Result
2002	16-Jul-2001	Spike in main field	17-Jul-2001	Double coil protection sheet
1027	29-Oct-2002	Missing outer shim	01-Nov-2002	Missing outer shim
3135	27-Jan-2004	Inward movement of block5 and 6	17-Feb-2004	Folded outer shim
1108	22-Apr-2004	Missing or additional thickenss on outer pole	12-Jul-2004	No visible defect
3254	06-Sep-2004	Inward movement of block5 and 6	14-Sep-2004	Folded outer shim
Bad quality of the coil gluing				
Magnet	Measured on	Analysis	Opened on	Result
2032	21-May-2003	Inward movement of block6	18-Nov-2003	Block6 detached from inner layer
2035	14-Jul-2003	Inward movement of block6	27-Apr-2004	Block6 detached from inner layer
1099	20-Feb-2004	Inward movement of block6	16-Mar-2004	Block6 detached from inner layer
3175	20-Apr-2004	Inward movement of block6	11-May-2004	Block6 detached from inner layer
1108	22-Apr-2004	Inward movement of block6	12-Jul-2004	Block6 detached from inner layer
1122	23-Apr-2004	Inward movement of block6	24-May-2004	Block6 detached from inner layer
1128	03-May-2004	Inward movement of block6	05-Jul-2004	Block6 detached from inner layer
1130	10-May-2004	Inward movement of block6	14-Jul-2004	Block6 detached from inner layer
Other				
Magnet	Measured on	Analysis	Opened on	Result
2065	15-Mar-2004	Inward movement of block6	29-Apr-2004	Good glue, movement observed
2089	18-May-2004	Inward movement of block6	01-Jun-2004	Good glue, no movement observed
2084	10-May-2004	Inward movement of block6	09-Jun-2004	Good glue, small movement observed

4.2 Estimated coil waviness

- Coil waviness estimated from the variation of the multipoles along the axis is in general below 30 microns. The recent part of the production is very stable, showing values of waviness below 25 microns.
- The crisis between collared coils 390th and 480th, due to inward radial movements of block6 in some spots along the magnet axis in Firm2 and Firm1, is over (see Fig. 33).




Fig. 33: Estimated coil waviness in the straight part of the measured collared coils (black dots: aperture 1, blue dots: aperture 2).

Acknowledgements

We wish to acknowledge all colleagues involved in the measurements at room temperature and at 1.9 K, and the Firm personnel involved in magnetic measurements. We thank R. Chamizo and P. Hagen for data validation and analysis. We finally acknowledge the project engineers and MTM-AS for support in the analysis.